

GROUP 4A INSECTICIDE

PULL HERE TO OPEN ►

 **Meridian[®]25WG**

syngenta[®]

Insecticide

For foliar and systemic control of insect pests in turf-grass, sod farms, interiorscape and landscape plants.

Active Ingredient:

Thiamethoxam ¹	25.0%
Other Ingredients:	75.0%
Total:	100.0%

¹ CAS No. 153719-23-4

Meridian[®] 25WG is a water-dispersible granule that contains 4 oz. of active ingredient per lb. of formulated product.

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

See additional precautionary statements and directions for use in booklet.

EPA Reg. 100-943 EPA Est. 67545-AZ-1

Product of India Formulated in the USA

SCP 943A-L3A 0410 318249

1 pound, 1 ounce (17 ounces)

Net Weight

TM

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
There is no specific antidote if Meridian 25WG is ingested. Induce emesis or lavage stomach. Treat symptomatically.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

FIRST AID (continued)
<p>HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p>

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

continued...

PRECAUTIONARY STATEMENTS (*continued*)

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

• Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address:

<http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

• Ground Water Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

• Spray Drift Advisory

Do not allow this product to drift.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material – Category A (e.g., natural rubber ≥ 14 mils)
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

USE INFORMATION

Meridian 25WG is a broad-spectrum insecticide that is effective when applied at label rates to turfgrass, plant foliage and soil.

Meridian 25WG is active against many sucking and chewing insect pests by contact and ingestion. Meridian 25WG is relatively short-lived on the surface of plant foliage, readily absorbed into plant tissues, and rainfast once it has dried. The rapid translaminar absorption and distribution within leaves provides excellent residual control of foliar feeding insects.

In the soil, the active ingredient in Meridian 25WG will control soil pests upon contact or ingestion and is also readily taken up by plant roots. The active ingredient moves upwards in the plant to the site of pest infestation.

Through feeding on the plant, pests are exposed to the active ingredient in Meridian 25WG. Feeding will stop within minutes to hours of exposure and be followed by death of the pest. The moderate persistence of Meridian 25WG in the soil and foliage also provides residual control of labeled pests. Meridian 25WG is not active as an ovicide or as an insect growth regulator. Because residues on leaf surfaces are quickly degraded, Meridian 25WG is compatible with beneficial arthropods.

Resistance Management

Meridian 25WG belongs to the neonicotinoid class of chemistry (Group 4A Insecticide). Pests which are resistant to insecticides in other chemical classes are not known to be cross-resistant to Meridian 25WG. However, insect pests are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of Meridian 25WG should conform to sound resistance management strategies established for the crop and use area. Strategies should include rotation of chemicals with different modes of action, limitation of the number of successive generations of the target pest being treated with the same compound and adherence to labeled directions.

Consult your local pest control advisor or extension office for additional methods for preventing resistance development. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insect pests on this label.

MIXING PROCEDURES

Prepare only the amount of spray mixture that is needed for the immediate application. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Meridian 25WG Alone

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Meridian 25WG to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Meridian 25WG has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Meridian 25WG + Tank Mixtures

Meridian 25WG is compatible with many other commonly used insecticides, fungicides, liquid fertilizers. Check compatibility before tank mixing.

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as Meridian 25WG, liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

Note: When using Meridian 25WG in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including Meridian 25WG. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using Meridian 25WG in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. Do not exceed any dosage rate, and follow the most restrictive label precautions and limitations. Do not mix this product with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility

Meridian 25WG is compatible with most insecticide, fungicide, and foliar nutrients products. However, test the physical compatibility of Meridian 25WG with tank-mix partners before use. To determine the physical compatibility of Meridian 25WG with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

FIRE ANTS - INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL PLANTS)

Application Rate

For control of fire ants (*Solenopsis* species) prepare a drench solution at a concentration of 1 - 3 ounces of Meridian 25WG per 10 gallons of water. Thoroughly mix solution and apply directly to mounds. For optimum control of small ant mounds (<6 inches in diameter at the surface) apply one gallon of the drench solution per mound. For optimum control of larger ant mounds, apply 2 - 3 gallons of the drench solution per mound. Direct the drench application at the center of the mound and include a 6 inch diameter circle around the center of the mound. Do not apply less than 0.5 gallon or more than 3 gallons of drench solution per mound.

Application Timing

For control of fire ants, make a single drench application when mounds appear to be active.

APPLICATION TO TURFGRASS

Sites of Application

Meridian 25WG can be applied to turfgrass on: golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields and sod farms.

Restrictions:

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply more than 17.0 ounces of Meridian 25WG (0.266 pounds thiamethoxam) per acre per year

Broadcast Applications to TO Turfgrass

Apply Meridian 25WG as a broadcast application to turfgrass for control of targeted pests. Use sufficient water volume (1.5 - 5 gal./ 1000 square feet) to uniformly distribute Meridian 25WG over the area being treated and to adequately move the active ingredient into the canopy and thatch layers.

To help prevent grubs, irrigate within seven (7) days of application to move Meridian 25WG into the root zone where grubs feed. To control grubs, irrigate within one (1) day of application to move Meridian 25WG into the root zone where grubs feed. Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may impact the performance of Meridian 25WG against white grubs and mole crickets. Do not mow turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into turfgrass.

Applications to Turfgrass

Use sites include turfgrass on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields, and sod farms.

Use Pattern	Pests	Application Timing	Dosage of Meridian 25WG
Broadcast	Larvae (white grubs) of: <i>Aphodius</i> species Asiatic garden beetle Billbugs (<i>Sphenophorus</i> species) Black turfgrass atanius European chafer Green June beetle Japanese beetle May or June beetle (<i>Phyllophaga</i> species) Northern masked chafer Oriental beetle Southern masked chafer	Apply up to 45 days before the historical peak of adult flight to 2nd instar grub of the species being targeted. For optimum control, treat from peak flight to peak egg hatch. To control grubs, make the application from egg hatch to second instar (grubs less than half their full size).	3.0 to 4.0 oz. per 10,000 sq.ft. or 12.7 to 17 oz. per acre
	Crane flies (<i>Tipula</i> species)	For optimum control, apply at oviposition (egg lay)	
	Mole crickets (suppression) (<i>Scapteriscus</i> species)	To suppress damage, treat from first egg hatch to peak egg hatch. See note below.	
	Chinch bugs (suppression) (<i>Blissus</i> species)	Apply when young nymphs are first observed. See note below.	
	Flea beetles Greenbugs Leafhoppers Sod webworms Spittlebugs	For optimum control, make application when populations are first observed.	

Use Pattern	Pests	Application Timing	Dosage of Meridian 25WG
Broadcast (continued)	Ants (excluding Carpenter, Fire, Harvester, and Pharaoh ants)	To control ants, treat when ant mounds are first observed. To achieve optimum control on green and tee surfaces, treat the affected area plus a surrounding 30 foot buffer. For fairways and roughs, treat at least 1.5 times the infested area. Water in to adequately move active ingredient to target area. For additional knockdown activity, utilize with Scimitar GC Insecticide in an insect control spray program. Consult the Scimitar GC Insecticide label for use.	3.0 to 4.0 oz. per 10,000 sq.ft. or 12.7 to 17 oz. per acre
Individual Fire Ant Mounds	Fire Ants (<i>Solenopsis</i> species) Refer to "FIRE ANTS - INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL PLANTS)" section of label.		

For specific information about developmental stages of the target pest and optimal timing of applications, consult with your State Cooperative Extension Service.

Note: For turf with heavy thatch (more than 0.75 inches), use the higher rates within the specified range. Meridian 25WG is not phytotoxic to any major turfgrass species.

Note: Meridian 25WG provides suppression of mole crickets and chinch bug populations on turfgrass. Suppression can mean either erratic control, ranging from good to poor, or a consistent level of control below that generally acceptable for commercial purposes.

APPLICATION TO INTERIOSCAPE AND LANDSCAPE ORNAMENTAL PLANTS

Sites of Application

Meridian 25WG can be applied to ornamentals grown in residential and commercial landscapes, parks, golf courses and interiorscapes.

Restriction:

- **DO NOT** exceed 17 ounces of Meridian 25WG (0.266 pounds Thiamethoxam) per acre per year.

Application Methods

Meridian 25WG is an effective insecticide when applied to plant foliage, soil or soil-less media, and fire ant mounds.

Foliar Application

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). On hard to wet plants, spray adjuvants that improve wetting of foliage are recommended. Do not use spray adjuvants that may bind Meridian 25WG to the leaf surface, which would limit absorption into the foliage. Meridian 25WG can be applied in a range of spray volumes (concentrate to dilute spray volumes) provided thorough and uniform coverage is obtained. For concentrate sprays (i.e., ULV, foggers, air assist equipment), apply the same amount of Meridian 25WG per unit area as would be used in a dilute spray volume over the same area.

Begin applications when labeled pests first appear or when economic thresholds have been reached. Repeat applications as necessary to maintain control, but no sooner than every 7 days.

Foliar Applications to Interiorscape and Landscape Ornamental Plants

Plant type	Pests	Dosage of Meridian 25WG	Application Timing
Trees	Ants (excluding Carpenter, Fire, Harvester, and Pharaoh ants)	2.0 to 4.0 oz. per 100 gal.	Apply when pests are first observed. Reapply as needed but no sooner than every 7 days.
Shrubs			
Evergreens	Aphids		
Flowers	Lace Bugs Leafhoppers Leaf Beetles		
Foliage Plants	Tent caterpillars	4.0 to 8.5 oz. per 100 gals.	For gall makers, apply during egg laying of the generation being targeted. For leafhoppers, plant bug, and aphids make the first application when populations are first observed. Repeat applications as necessary to maintain control, but no sooner than 7 days after the last application.
Groundcovers	Adelgids (including Hemlock woolly adelgid)		
Interior Plantscapes	Flea Beetles		
	Japanese beetle adults		
	Leaf gall forming insects		
	Leafminers		
	Mealybugs		
	Midges		
	Plant Bug		
	Psyllids (including Asian Citrus Psyllid)		
	Sawflies		
Soft Scales	For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes.		
Spittlebugs			
Thrips- foliar feeding (suppression)			
	Whiteflies		

continued...

Foliar Applications to Interiorscape and Landscape Ornamental Plants (continued)

Plant type	Pests	Dosage of Meridian 25WG	Application Timing
Trees Shrubs Evergreens Flowers Foliage Plants Groundcovers Interior Plantscapes	Adult Black vine weevil (suppression)	4.0 oz. per 50 gal. or 8.0 oz. per 100 gal.	Apply to foliage of plants when adult feeding damage is first observed. Make only one application per generation. For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes. For best control, use in a rotational spray program with Scimitar CS Insecticide. Consult Scimitar CS Insecticide label for use.

For specific information about developmental stages of the target pest, associated damage and action thresholds to properly time applications consult with your State Cooperative Extension Service.

Application to Soil and Soil-less Media of Ornamental Plants

Meridian 25WG is effective as a soil and systemic insecticide when applied to the soil or soil-less media using broadcast, soil injection, or soil drench applications.

As a broadcast application, use sufficient water volume (1.5 – 5 gallons/1000 square feet) to uniformly distribute Meridian 25WG at the labeled rate over the area being treated.

For soil injection around trees and shrubs, uniformly select sites and evenly distribute Meridian 25WG solution among injection sites.

For soil drench applications around the base of trees and shrubs, apply the proper rate uniformly in a minimum of 16 fluid ounces of solution per inch diameter of breast height (D.B.H.) or per foot of shrub height.

Meridian 25WG must move to the feeding site of the target pest, whether this is in the root zone or the above ground portion of the host plant. Therefore, control of pests from the systemic activity of Meridian 25WG may be delayed for one or more weeks depending on plant size, rate of uptake from the soil or rate of translocation within the plant. Systemic activity in large plants may require several weeks before control of stem and foliar pests is achieved. If economic thresholds for target pests are low, or plants are historically damaged, apply Meridian 25WG preventatively (before pests are evident) to the growing media.

Irrigate within 3 days of treatment (optimally within 24 hours) to move Meridian 25 WG into the root zone of plants or to the location of soil-dwelling pests. Do not over irrigate during the first seven days after treatment or reduced efficacy may occur.

Applications to Soil of Ornamentals Plants

For plants grown in residential and commercial landscapes, parks, golf courses and interiorscapes.

A. Soil Inhabiting Pests

Plant Types	Pests	Dosage of Meridian 25WG	Application Timing
All plant types	Larvae (Grubs) of:	12.7 to 17 oz. per acre or 3.0 to 4.0 oz. per 10,000 sq. ft.	Application Timing for Grubs: Apply up to 45 days before the historical peak of adult flight to 2 nd instar grub of the species being targeted. For optimum control, treat from peak flight to peak egg hatch. To control grubs, make the application from egg hatch to second instar (grubs less than half their full size).
	<i>Aphodius</i> species		
	Asiatic garden beetle		
	Black turfgrass ataenius		
	European chafer		
	Green June beetle		
	Japanese beetle		
	May or June beetle (<i>Phyllophaga</i> species)		
	Northern masked chafer		
	Oriental beetle		
Southern masked chafer			

Plant Types	Pests	Dosage of Meridian 25WG	Application Timing
All plant types (continued)	Fire Ants (<i>Solenopsis</i> species) Refer to "FIRE ANTS - INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL PLANTS)" section of label.		

- **Note:** If leaching, wait a minimum of 7 days after application to allow for maximum uptake of Meridian 25WG by the plant's root system.
- Apply to the soil preventively in situations where tolerance of insects or damage is low. Allow a minimum of one week for smaller plants and at least 2 weeks for large plants to translocate Meridian 25WG to feeding sites of the target pest.
- Irrigate after application for optimal performance.
- **Do not** exceed 17 ounces/Acre per year when making applications to the soil for ornamentals.
- For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service. Refer to additional comments under **Use Information** section relating to ornamental applications to soil and related irrigation events.

B. Above Ground Pests

Pests	Plant Types	Dosage of Meridian 25WG	Application Instructions
Adelgids Aphids Lace Bugs Flea beetles Japanese beetle adults Leaf beetles Leafminers Mealybugs Root aphids Root Weevil larvae (including <i>Diaprepes abbreviatus</i>) Sawflies Soft Scales Spittlebugs Tent caterpillars Thrips-foliar feeding (suppression) Whiteflies	Trees	Single Tree 0.07 to 0.14 oz. (2 to 4 grams) per one inch of tree trunk diameter (D.B.H.)	Soil Injection Method For Trees <i>Basal System:</i> Apply the proper rate among 4 evenly spaced injection sites located within a foot of each trunk. <i>Grid System:</i> Apply the proper rate among injection sites on a grid pattern with 1.5-2.5 feet centers extending to the drip line of the tree. Soil Drench: Apply the proper rate uniformly around the base of the tree within 1-2 feet of the trunk. Drench targeted area with the Meridian 25 WG solution without runoff. Use a minimum of 16 fluid ounces solution per inch D.B.H. Irrigate after application for optimal performance.

Pests	Plant Types	Dosage of Meridian 25WG	Application Instructions
Adelgids Aphids Lace Bugs Flea beetles Japanese beetle adults Leaf beetles Leafminers Mealybugs Root aphids Root Weevil larvae (including <i>Diaprepes abbreviatus</i>) Sawflies Soft Scales Spittlebugs Tent caterpillars Thrips-foliar feeding (suppression) Whiteflies	Shrubs	Single Shrub 0.07 to 0.14 oz. (2 to 4 grams) per foot of shrub height Hedge Row Apply 0.07 to 0.14 ounces (2 to 4 grams) per foot of shrub height (<6' tall) per plant	Soil Injection Method For Shrubs Within the canopy, apply the proper rate among a minimum of 4 evenly spaced injection sites per shrub. For multiple plants (hedge row), determine the number of plants per hedge row, multiply by the average height and rate to be applied per foot of height. Soil Drench: Apply the proper rate uniformly around the shrub within 1-2 feet of the base. Drench targeted area with the Meridian solution without runoff. Use a minimum of 16 oz solution per foot shrub height. For multiple plants (hedge row), determine the number of plants per hedge row, multiply by the average height and rate to be applied per foot of height. Irrigate after application for optimal performance.

B. Above Ground Pests (continued)

Pests	Plant Types	Dosage of Meridian 25WG	Application Instructions
Adelgids Aphids Lace Bugs Flea beetles Japanese beetle adults Leaf beetles Leafminers Mealybugs Root aphids Root Weevil larvae (including <i>Diaprepes abbreviatus</i>) Sawflies Soft Scales Spittlebugs Tent caterpillars Thrips-foliar feeding (suppression) Whiteflies	Annuals & small woody landscape plants	3.0 to 4.0 oz. per 10,000 sq. feet	Apply as a broadcast treatment to the soil directed to the plant root zone. Irrigate after application for optimal performance.

Application Timing for Above Ground Pests:

To Help Prevent Pests: Apply 1 to 4 weeks before pest has historically been observed.

To Control Pests: Allow 1 to 4 weeks for reduction in pest density to occur. **Note:** Larger plants require the longer interval.

- **Note:** If leaching, wait a minimum of 7 days after application to allow for maximum uptake of Meridian 25WG by the plant's root system.
- Apply to the soil preventively in situations where tolerance of insects or damage is low. Allow a minimum of one week for smaller plants and at least 2 weeks for large plants to translocate Meridian 25WG to feeding sites of the target pest.
- **Do not** exceed 17 ounces/Acre per year when making applications to the soil for ornamentals.

For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service. Refer to additional comments under **Use Information** section relating to ornamental applications to soil and related irrigation events.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

Pesticide Disposal



Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (plastic containers)

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling (bag)

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Meridian®, Scimitar, the ALLIANCE FRAME 
the SYNGENTA Logo and the PURPOSE ICON 
are trademarks of a Syngenta Group Company
©2010 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 943A-L3A 0410
318249

GROUP 4A INSECTICIDE



Insecticide

For foliar and systemic control of insect pests in turfgrass, sod farms, interiorscape and landscape plants.

Active Ingredient:	
Thiamethoxam ¹	25.0%
Other Ingredients:	75.0%
Total:	100.0%

¹CAS No. 153719-23-4

Meridian® 25WG is a water-dispersible granule that contains 4 oz. of active ingredient per lb. of formulated product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. 100-943
EPA Est. 67545-AZ-1

Meridian® and the Syngenta logo are trademarks of a Syngenta Group Company
©2010 Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

SCP 943A-L3A 0410 318249

1 pound, 1 ounce
(17 ounces) Net Weight

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements and directions for use in booklet.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID: If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: There is no specific antidote if Meridian 25WG is ingested. Induce emesis or lavage stomach. Treat symptomatically. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372.

Environmental Hazards: This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

syngenta®

BAR CODE # IS
(01) 0 07 02941 35036
LAST DIGIT IS CHECK DIGIT
UCC/EAN 128

KEEP OUT OF REACH OF CHILDREN CAUTION

- **Surface Water Advisory**

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

- **Ground Water Advisory**

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

- **Spray Drift Advisory**

Do not allow this product to drift.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¹/₄ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Chemigation: DO NOT apply this product through any type of irrigation system.

Meridian® and the Syngenta logo are trademarks of a Syngenta Group Company
©2010 Syngenta

Manufactured for:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 943A-L4A 0410 318250